

# Topological Description of the Aging Dynamics in Simple Glasses

L. Angelani<sup>1</sup>, R. Di Leonardo<sup>2</sup>, G. Parisi<sup>3</sup>, and G. Ruocco<sup>3</sup>

<sup>1</sup> *Università di Trento and INFM, I-38050, Povo, Trento, Italy.*

<sup>2</sup> *Università di L'Aquila and INFM, I-67100, L'Aquila, Italy.*

<sup>3</sup> *Università di Roma La Sapienza and INFM, I-00185, Roma, Italy.*

(February 1, 2008)

We numerically investigate the aging dynamics of a monatomic Lennard-Jones glass, focusing on the topology of the potential energy landscape which, to this aim, has been partitioned in basins of attraction of stationary points (saddles and minima). The analysis of the stationary points visited during the aging dynamics shows the existence of two distinct regimes: *i*) at short times,  $t < t_c$ , the system visits basins of saddles whose energies and orders decrease with  $t$ ; *ii*) at long times,  $t > t_c$ , the system mainly lies in basins pertaining to minima of slowly decreasing energy. The dynamics for  $t > t_c$  can be represented by a simple random walk on a network of minima with a jump probability proportional to the inverse of the waiting time.

PACS Numbers : 61.20.Lc, 64.70.Pf, 02.70.Ns

The very slow approach to equilibrium of glassy systems is a very interesting and intriguing physical process that, in the last few years, has received considerable attention from analytical, numerical and experimental point of view [1]. When a glassy system is suddenly brought in the glassy state from an high temperature (or low density) equilibrium state, it starts to explore the phase space with an off-equilibrium dynamics which slows down as the system “ages”. During this aging regime one time physical quantities, that are time independent on average at equilibrium, slowly approach their equilibrium values, while two time quantities, like the correlation functions, depend on both times due to the lack of time translation invariance. The study of the off-equilibrium dynamics of glassy systems, and in particular of spin-glasses, has allowed to obtain important informations on the phase-space properties of the systems itself (behavior of the order parameter in the low temperature equilibrium phase [2–5]) and to formulate off-equilibrium relations generalizing the usual equilibrium ones (generalized fluctuation-dissipation relation [1,2]) with the introduction of an effective temperature [6]. The investigation of the off-equilibrium properties seems to indicate the correctness of the conjecture stating the similarity of structural glasses with some spin glass model (spin-glasses with one step replica symmetry breaking) [7–11]. The way a glassy system goes towards equilibrium is then not only an interesting problem by itself but it is also of great relevance in view of a deeper understanding of the nature of glass transition. In this respect, many efforts have been devoted to the understanding of the role played by the potential energy landscape in the equilibrium dynamics of supercooled liquids [12]. The trajectory of the representative point in the  $3N$  configuration space can be mapped into a sequence of locally stable points (the so called *Inherent Structures*, IS [13]), that are the local minima of the total potential energy  $V$ : to each instantaneous configuration during the dynamical evolution of the system one can associate an IS by a

steepest descent path in the  $V$  surface. The properties of the IS has been found to be very useful to clarify many features in the dynamics and the thermodynamics of supercooled liquids in [14–17] and off [18,19] equilibrium. An extension of such an analysis including all stationary points of  $V$  (minima and saddles) [20–22] has been recently proposed to form a better representation of the system during the equilibrium dynamics. This approach gives a novel point of view on the equilibrium dynamics of supercooled liquids based on a detailed topological analysis of the potential energy landscape. In particular it emerges [21] that the order of the sampled saddles during the equilibrium dynamics is a well defined increasing function of temperature and it appears to vanish at the so-called mode-coupling [25] transition temperature  $T_{MCT}$ , which then marks the crossover from a dynamics between basins of saddles ( $T > T_{MCT}$ ) to a dynamics between basins of minima ( $T < T_{MCT}$ ). Similar results have been found [23] in the framework of Instantaneous Normal Modes analysis [24].

In this Letter we numerically investigate the off-equilibrium dynamics of a simple model glass, by mapping the true dynamics in the configuration space on to the stationary points of  $V$ . In our glass the particles interact via a Lennard-Jones potential modified in such a way as to avoid crystallization [26]. The system is brought in an off-equilibrium initial condition by a sudden isothermal density jump (*crunch*) across the liquid-glass transition line. We find a rather sharp cross-over between two dynamical regimes: *i*) a short times regime where the system point travels between basins of saddles of decreasing energy and order and *ii*) a long time regime where the system point moves between basins of saddles of zero order (minima) and decreasing energy, crossing basins of saddles of low order. The existence of two well separated dynamical regimes shows up also in the Mean Square Displacement (MSD). Here a logarithmic behavior is found at long times suggesting the picture of a system point performing a random walk between basins of

minima with a jump probability per unit time inversely proportional to time elapsed since crunch. This picture is quantitatively supported by the very good agreement found between the MSD measured along the MD trajectories and the MSD reconstructed *via* a random walk model.

The system under study is a monatomic Modified-Lennard-Jones (MLJ) [26] that is the usual 6-12 Lennard-Jones pair interaction (truncated and shifted at  $R_c = 2.6$ , standard Lennard-Jones units are used hereafter) plus a small many body term which inhibits crystallization with negligible corrections to the equation of state of a monatomic Lennard-Jones system (see [11,26] for details). The system is composed of  $N = 256$  particles enclosed in a cubic box with periodic boundary conditions. The initial liquid equilibrium configurations ( $\rho_0 = 0.95$ ,  $T_0 = 0.5$ ) are prepared by standard isothermal MD. At time  $t = 0$  the density is suddenly increased to the value  $\rho_1 = 1.24$ , where the glass transition temperature ( $T_g(\rho_1) \sim 1.4$  [11]) is well above the simulation temperature  $T_0$ . The subsequent evolution of the system is generated up to time  $t = 3 \cdot 10^4$  by means of standard microcanonical MD with time step  $\tau = 0.01$ . For each MD trajectory we quenched a set of logarithmically equally spaced configurations to the corresponding IS and saddle following the steepest descent path respectively on the potential energy surface  $V$  in the former case and on the auxiliary potential  $W = \frac{1}{2}|\vec{\nabla}V|^2$  [20–22] in the latter.




FIG. 1. Energies as a function of time during off equilibrium dynamics. Instantaneous configuration potential energy per particle  $e_i$  subtracted by the constant term  $\frac{3}{2}T_0$  (open circles), inherent structure saddle energy  $e_s$  (full triangle) and the inherent structure minimum energy  $e_m$  (open diamonds). Error bars are shown only when larger than symbol size. The arrow indicates time  $t_c$  where the crossover from dynamics between basins of saddles to dynamics between basins of minima takes place.

In Fig.1 we report in a logarithmic scale the time evolution of: a) the instantaneous configuration potential energy per particle  $e_i$  subtracted of the constant term  $\frac{3}{2}T_0$  (open circles), b) the saddle's energy  $e_s$  (full triangles), c)

the IS energy  $e_m$  (open diamonds). The remarkable coincidence between  $e_s$  and  $e_i - \frac{3}{2}T_0$  suggests to think of the instantaneous energy  $e_i$  as the sum of two contributions: the saddle's energy (slowly approaching equilibrium) and the energy of nearly  $N$  fast harmonic equilibrated degrees of freedom. It is evident the presence of a rather sharp transition in dynamical behavior of  $e_i$  and  $e_s$  at time  $t_c \sim 20$ . In particular the time evolution of  $e_i$  for  $t > t_c$  seems to be driven by the inherent dynamics between minima as suggested by the underlying time evolution of  $e_m$ .

In Fig.2 the mean saddle's order  $n_s$  (number of negative eigenvalues of the potential energy Hessian computed on the saddle configurations) is reported as a function of time. The dynamic transition observed at  $t_c$  in the energy time evolution in Fig.1 corresponds here in the mean order  $n_s$  falling below the value  $n_s = 1$  or, in other words, for times longer than  $t_c$  the system point starts to explore extensively the basins of saddles of order zero (minima). The inset in Fig.2 shows the order  $n_s$  computed along a single MD trajectory. One can see that for times  $t > t_c$  the systems visits basins of minima passing through basins of low order saddles.




FIG. 2. Mean inherent saddle order  $n_s$  (number of negative eigenvalues of the Hessian matrix of potential energy evaluated at the inherent structure saddle) as a function of time  $t$  after crunch. The inset shows  $n_s$  computed along a single trajectory.

In a recent study of the MLJ landscape probed in equilibrium [21] some of us found that the mean elevation energy of a saddle of order  $n_s$  from the underlying minimum is proportional to  $n_s$  (this is at variance with the result reported in [22], where a linear relation is observed between  $n_s$  and the *absolute* energy of the saddles). This is clearly a property of the energy landscape itself and one expects that the same properties are also valid in the off-equilibrium dynamics.

In Fig.3 where we report  $e_s$  together with  $e_m + n_s \Delta E/N$  with  $\Delta E$  a fit parameter. This result confirms the picture of an energy landscape where the saddles of

order  $n_s + 1$  lay above the saddles of order  $n_s$  of a constant quantity  $\Delta E$ . From the data we obtain the value  $\Delta E = 10$ . The value of  $\Delta E$  could also be obtained from an equilibrium analysis, as in [21] where a value of  $\Delta E = 3.6$  is found for the case  $\rho = 1$ . We have not carried out such an equilibrium analysis for the density  $\rho = 1.24$  used here.




FIG. 3. Saddle's energy  $e_s$  (solid symbols) evaluated during off equilibrium dynamics compared with the energy of the minima  $e_m$  increased by the quantity  $n_s \Delta E / N$  (open symbols) where  $n_s$  is the time dependent saddle's order and  $\Delta E = 10$  is a fitting constant representing the mean energy separation between a saddle of order  $n_s + 1$  and a saddle of order  $n_s$ .




FIG. 4. Mean square displacement  $r^2(t, t_w)$  as a function of time  $t$  for four selected  $t_w$  values: 1, 5, 20 and 200 (solid symbols). The straight lines are the prediction of a random walk model between minima with jump probability per unit time decreasing as the inverse of time,  $p(t) = c/t$ . According to the model the slope of these lines are  $d_0^2 c$  where  $d_0^2$  is the mean square distance between adjacent minima (see text below for details). The arrow indicates time  $t_c$  where the crossover from dynamics between basins of saddles to dynamics between basins of minima takes place. The system size is  $N = 256$ .

Another interesting quantity in describing the dynamical processes is the mean square displacement of the configuration at time  $t$  from that at time  $t_w$  after crunch:

$$r^2(t, t_w) = \frac{1}{N} \sum_i \langle |\mathbf{r}_i(t) - \mathbf{r}_i(t_w)|^2 \rangle, \quad (1)$$

where  $\langle \dots \rangle$  is an average over initial configurations. In Fig.4 we report  $r^2(t, t_w)$  as a function of time  $t$  for four selected  $t_w$  values:  $t_w = 1, 5, 20, 200$ . Also in this case  $t_c \sim 20$  marks a transition in the dynamical behavior. In particular the long time regime shows a well defined logarithmic dependence on time. This suggests the picture of the diffusion process proceeding through elementary uncorrelated events occurring with a probability per unit time  $p(t) \propto 1/t$  [27]. As previously observed, in the long time regime the system point travels between basins of minima. It is therefore natural to identify the elementary diffusion events as jumps between adjacent basins of minima. In order to give further support to this conjecture we calculated the cumulative number of jumps  $N_{jumps}(t)$  defined as the number of jump events in which the constantly monitored energy of the IS makes a sharp transition.




FIG. 5. a) Cumulative number of jumps between adjacent minima visited by the system during the off-equilibrium dynamics; the solid line is the best fit for  $t > 20$  with slope  $c = 1.0$ . b) Mean square distance between adjacent minima  $d^2$ ; the solid line is the constant value  $d_0^2 \sim 0.019$  reached for times greater than  $t_c \sim 20$ . The system size is  $N = 256$ .

The quantity  $N_{jumps}$  is plotted in Fig.5(a). For times  $t > t_c$   $N_{jumps}$  displays a nice logarithmic behavior that

is what one would expect if the probability per unit time of a jump event is  $p(t) = c/t$ . In other words the time needed to escape from an IS reached by the system after a time  $t_w$  from *crunch* is proportional to time  $t_w$  itself (a similar behavior is found in the off-equilibrium dynamics of a ferromagnetic Ising system where the persistence time of a bubble in which spins have a similar value is proportional to time  $t_w$  [28,29]). In Fig.5(a) the solid line represents the fitting function  $c \log(t)$  with  $c = 1.0$ . In Fig.5(b) we report the mean square distance  $d^2$  between adjacent minima probed by the system during the aging dynamics:

$$d^2(t) = \frac{1}{N} \sum_i \langle |\mathbf{r}_i^{IS}(t^+) - \mathbf{r}_i^{IS}(t^-)|^2 \rangle, \quad (2)$$

where  $t$  is the time of the jump between two different basins of minima and  $\mathbf{r}_i^{IS}(t^-)$ ,  $\mathbf{r}_i^{IS}(t^+)$  are the coordinates of IS respectively before and after time  $t$ . The quantity reported in figure approaches the long time constant value  $d_0^2 \sim 0.019$ . For uncorrelated jumps, the mean square displacement would be simply given by

$$r^2(t, t_w) = d_0^2 c \log(t/t_w). \quad (3)$$

This is in a remarkably good agreement with the MD data as shown in Fig.4, where we draw straight lines with slope  $d_0^2 c = 0.019$  (as obtained from the values of  $d_0^2$  and  $c$  derived from Fig.5) and intercept chosen as to best fit the data for  $t > 20$ .

In conclusion we have numerically studied the off-equilibrium dynamics of a monatomic Modified-Lennard-Jones system by sudden isothermal density jumps from the liquid to the glassy phase. Analyzing the time evolution of the energy and the mean square displacement we found two well defined dynamical regimes, which can be related to the different ways the system explores the potential energy landscape: *i*) at short times the system point moves downward passing through basins of saddles with decreasing energy and order, and *ii*) at longer time the relevant dynamical processes are jumps between basins of minima with slower decreasing energy. In the whole time regime the instantaneous potential energy  $e_i$  can be thought as due to a sum of two contributions: the energy of the underlying saddle's configurations  $e_s$  plus a nearly harmonic term of the fast degrees of freedom  $3/2T$ . Moreover the linear energy gap between saddles and underlying minima  $e_s - e_m$  as a function of saddles order  $n_s$  supports the picture of a simple energy landscape with only a single energy barrier parameter  $\Delta E$  which represents the energy gap between a saddle of order  $n_s$  and a saddle of order  $n_s + 1$ . The relevant processes in the long time aging regime are well represented by a simple random walk model with elementary events corresponding to jumps between minima with a probability per unit time proportional to the inverse of time elapsed since the system has been brought out of equilibrium.

- 
- [1] For a review see J.P. Bouchaud, L.F. Cugliandolo, J. Kurchan and M. Mezard, *Out of equilibrium dynamics in spin glasses and the other glassy systems*, in *Spin Glasses and Random Fields*, edited by P. Young, World Scientific (1997).
  - [2] L.F. Cugliandolo and J. Kurchan, Phys. Rev. Lett. **71**, 173 (1993).
  - [3] S. Franz and M. Mézard 1994, Europhys. Lett. **26**, 209.
  - [4] S. Franz, M. Mézard, G. Parisi, L. Peliti, Phys. Rev. Lett. **81**, 1758 (1998).
  - [5] E. Marinari, G. Parisi, F. Ricci-Tersenghi, J. Ruiz-Lorenzo and F. Zuliani, J. Stat. Phys. **98**, 973 (2000).
  - [6] L.F. Cugliandolo, J. Kurchan and L. Peliti Phys. Rev. E **55**, 3898 (1997).
  - [7] T.R. Kirkpatrick and P.G. Wolynes, Phys. Rev. B **35**, 3072 (1987); T.R. Kirkpatrick and D. Thirumalai, Phys. Rev. B **36**, 5388 (1987); T.R. Kirkpatrick and D. Thirumalai, Phys. Rev. Lett. **58**, 2091 (1987).
  - [8] S. Franz and G. Parisi, Phys. Rev. Lett. **79**, 2486 (1997); M. Mézard and G. Parisi, Phys. Rev. Lett. **82**, 747 (1998); B. Coluzzi, G. Parisi and P. Verrocchio Phys. Rev. Lett. **84**, 306 (2000).
  - [9] G. Parisi, Phys. Rev. Lett. **79**, 3660 (1997).
  - [10] J.L. Barrat and W. Kob, Europhys. Lett. **46**, 637 (1999).
  - [11] R. Di Leonardo, L. Angelani, G. Parisi, and G. Ruocco, Phys. Rev. Lett. **84**, 6054 (2000).
  - [12] Special issue of Physica (Amsterdam) **107D**, Issue: 2-4 (1997).
  - [13] F.H. Stillinger and T.A. Weber, Phys. Rev. A **25**, 978 (1982); Science **225**, 983 (1984); F.H. Stillinger, Science **267**, 1935 (1995).
  - [14] S. Sastry, P.G. Debenedetti, and F.H. Stillinger, Nature **393**, 554 (1998).
  - [15] S. Buechner and A. Heuer, Phys. Rev. E **60**, 6507 (1999).
  - [16] F. Sciortino, W. Kob, and P. Tartaglia, Phys. Rev. Lett. **83**, 3214 (1999).
  - [17] T.B. Schröder, S. Sastry, J. Dyre, and S.C. Glotzer, J. Chem. Phys. **112**, 9834 (2000).
  - [18] W. Kob, F. Sciortino and P. Tartaglia, Europhys. Lett. **49**, 590 (2000).
  - [19] F. Sciortino and P. Tartaglia, cond-mat/0007208.
  - [20] A. Cavagna, cond-mat/9910244.
  - [21] L. Angelani, R. Di Leonardo, G. Ruocco, A. Scala, and F. Sciortino, Phys. Rev. Lett. (to be published).
  - [22] K. Broderix, K.K. Batthacharya, A. Cavagna, A. Zippelius, and I. Giardina, Phys. Rev. Lett. (to be published).
  - [23] C. Donati, F. Sciortino, and P. Tartaglia, Phys. Rev. Lett. **85**, 1464 (2000).
  - [24] T. Keyes, J. Phys. Chem. **101**, 2921 (1997).
  - [25] W. Götze, J. Phys.: Condens. Matter **11**, A1 (1999).
  - [26] G. Ruocco, L. Angelani, and R. Di Leonardo (to be published).
  - [27] G. Parisi, Phys. Rev. Lett. **79**, 3660 (1997).
  - [28] G. Parisi, J. Phys. Chem. B **103**(20), 4128 (1999).
  - [29] A.J. Bray, Adv. Phys., **43**, 357 (1994).